



US005810985A

# United States Patent [19]

[11] Patent Number: **5,810,985**

Bao et al.

[45] Date of Patent: **Sep. 22, 1998**

[54] **ELECTROPHORETICALLY MEDIATED CHEMICAL ANALYSIS**

[75] Inventors: **Jianmin Bao; Bryan Joseph Harmon; Dale Hansen Patterson; Fred E. Regnier**, all of West Lafayette, Ind.

[73] Assignee: **Purdue Research Foundation**, West Lafayette, Ind.

[21] Appl. No.: **386,224**

[22] Filed: **Feb. 9, 1995**

### Related U.S. Application Data

[63] Continuation of Ser. No. 944,846, Sep. 14, 1992, abandoned.

[51] Int. Cl.<sup>6</sup> ..... **C25B 7/00**; G01N 33/573

[52] U.S. Cl. .... **204/451**; 204/452; 204/601; 204/603; 204/182.1; 204/180.1; 204/182.8; 204/299 R; 435/7.4

[58] Field of Search ..... 204/182.1, 182.8, 204/299 R, 451, 452, 601, 603; 435/7.4

[56] **References Cited**

#### U.S. PATENT DOCUMENTS

4,181,589	1/1980	Brooks	.....	204/180 R
4,312,727	1/1982	Shainoff	.....	204/180 G
4,756,884	7/1988	Hillman et al.	.....	435/810
4,865,706	9/1989	Karger et al.	.....	204/182.8
4,865,707	9/1989	Karger et al.	.....	204/182.8
5,045,172	9/1991	Guzman	.....	204/299 R
5,061,361	10/1991	Gordon	.....	204/299 R
5,084,150	1/1992	Karger et al.	.....	204/180.1
5,085,756	2/1992	Swedberg	.....	204/299 R
5,089,099	2/1992	Chien et al.	.....	204/180.1
5,116,471	5/1992	Chien et al.	.....	204/180.1
5,122,248	6/1992	Karger et al.	.....	204/182.8
5,137,609	8/1992	Manian et al.	.....	204/180.1
5,145,567	9/1992	Hsieh et al.	.....	204/108.1
5,264,095	11/1993	Hsieh	.....	204/180.1

#### FOREIGN PATENT DOCUMENTS

2 040 313	8/1980	United Kingdom .
WO 91/06850	5/1991	WIPO .

#### OTHER PUBLICATIONS

Aebersold et al., Analysis of Dilute Peptide Samples by Capillary Zone Electrophoresis, 1990, J. Chromatogr., 516:79-88.

Chien et al., Field Amplified Sample Injection in High-Performance Capillary Electrophoresis, 1991, J. of Chromatogr., 559:141-152.

Nashabeh et al., Capillary Zone Electrophoresis of Proteins with Hydrophilic Fused-Silica Capillaries, 1991, Chromatogr., 599:367-383.

Emmer et al., Improved Capillary Zone Electrophoretic Separation of Basic Proteins, Using a Fluorosurfactant Buffer Additive, 1991, J. Chromatogr., 547:544-550.

Bruin et al., Capillary Zone Electrophoretic Separations of Proteins in Polyethylene Glycol-Modified Capillaries, 1989, 471:429-436.

Jorgenson et al., Zone Electrophoresis in Open-Tubular Glass Capillaries, 1981, Analytical Chemistry, , 53:1298-550.

Lauer et al., Capillary Zone Electrophoresis of Proteins in Untreated Fused Silica Tubing, 1986, Anal. Chem., 58:166-170.

Nielsen et al., Capillary Zone Electrophoresis of Insulin and Growth Hormone, 1989, Anal. Biochem., 177:20-26.

Schlabach et al., Rapid Assessment of Isoenzymes by High-Performance Liquid Chromatography, 1978, Clinical Chemistry, 24:1351-1360.

Snyder, Reaction Colorimeters as Detectors in High-Performance Liquid Chromatography, 1976, Journal of Chromatography, 125:287-306.

Towns et al., Synthesis and Evaluation of Epoxy Polymer Coatings for the Analysis of Proteins by Capillary Zone Electrophoresis, 1992, J. Chromatogr., 599:227-237.

Wu et al., High-Performance Capillary Isoelectric Focusing with a Concentration Gradient Detector, 1992, Anal. Chem., 64:219-224.

Yoshida, Hemolytic Anemia and G6PD Deficiency, 1973, Science, 179:532-537.

Primary Examiner—Cecilia J. Tsang

Assistant Examiner—C. Delacroix-Muirheid

Attorney, Agent, or Firm—Testa, Hurwitz & Thibault, LLP

[57] **ABSTRACT**

Disclosed are methods of electrochemical analysis of an analyte in a sample. The methods include providing a sample comprising an analyte, and introducing the sample into a capillary tube containing electrophoretic running buffer and a reactant. According to the methods of the invention, one or both of the analyte and the reactant is electrically charged. Chemical contact between analyte and reactant induced by their relative electrophoretic mobility results in the breaking or formation of a covalent bond of one of the analyte or the reactant to produce a detectable product. The methods also include the steps of imposing along the length of the capillary tube an electric current for a time sufficient to bring into chemical contact within the tube analyte and reactant so as to allow formation of product, and detecting the product.

**5 Claims, 17 Drawing Sheets**